

New Mexico Childcare/Pre-School/School Entry Immunization Requirements: 2022-23 school year

New Mexico School Nurses are granted Public Health authority by the NM Secretary of Health for collecting and submitting immunization information

Vaccine	Minimum # of vaccine doses by childcare and pre-school age levels							Vaccine doses by school grade level												Notes		
	By 4 Mo.	By 6 Mo.	By 12 Mo.	By 15 Mo.	16-47 Mo.	48-59 Mo.	≥ 60 Mo.	K	1 st	2 nd	3 rd	4 th	5 th	6 th	7 th	8 th	9 th	10 th	11 th		12 th	
Diphtheria/Tetanus/Pertussis (DTaP/DT/Td)*	1	2	3	3	3	4	5 (4)	5 (4)	5 (4)	5 (4)	5 (4)	5 (4)	5 (4)	5 (4)	5 (4)	5 (4)	5 (4)	5 (4)	5 (4)	5 (4)	5 (4)	One dose required on/after 4 th birthday. Four doses are sufficient if last dose given on/after 4 th birthday, with at least 6 months between the last two doses. Five doses are preferred for optimal protection.
Tetanus/Diphtheria/Pertussis (Tdap)																1	1	1	1	1	1	One dose Tdap required for entry into 7 th - 12 th grade.
Polio (IPV)* (OPV ²)	1	2	2	2	3	4 (3)	4(3)	4	4	4	4	4	4	4	4	4	4	4	4	4	4	Students in K-11 th grades final dose required on or after 4 th birthday. Three doses sufficient if CDC's catch-up schedule used AND last dose was given on/after 4 th birthday with at least 6 months between the last two doses.
Measles/Mumps/Rubella (MMR)				1	1	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	Min. age for valid 1 st dose is 12 months. Live vaccines (MMR, Varicella) must be given on the same day; if not, they must be administered a min. of 28 days apart.
Haemophilus Influenzae type B (Hib)*	1	2	2	2	3/2/1	3/2/1	3/2/1															
Hepatitis B (HepB)	1	2	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	Two doses adult Recombivax HB is also valid if administered at ages 11-15 years and if dose 2 received no sooner than 16 weeks after dose 1.
Pneumococcal (PCV)*	2	3	3	4/3/2/1	4/3/2/1	4/3/2/1	4/3/2/1/0															
Varicella (VAR)				1	1	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	Min. age for 1 st dose is 12 mos. Dose 2 should ideally be given at age ≥ 4, see note below. Live vaccines (MMR, Varicella) must be given on the same day; if not, they must be administered a minimum of 28 days apart.
Hepatitis A (HepA)				1	1	2	2	2	2													Hep A Vaccine is recommended for catch up in grades 2-12.
Meningococcal Men ACWY															1	1	1	1				1 dose required for 7 th -10 th grade entry. Booster dose recommended at age 16-18. Men ACWY strongly recommended for catch up for grades 11-12.

Diphtheria/Tetanus/Pertussis: If child (4 months-6 years) is behind schedule, follow the CDC's catch-up schedule.

Tetanus/Diphtheria/Pertussis: 7th-12th graders require proof of 1 dose of Tdap regardless of when the last Td-containing vaccine was given.

Catch-up: Children 7-18 years who are not fully immunized with the childhood DTaP series should be vaccinated according to the CDC's catch-up schedule, with Tdap as the 1st dose followed by Td if needed. A 3-dose series is sufficient if initiated after age 7, in which one dose must be Tdap, followed by 2 doses of Td. Children age 7-10 who receive 1 dose Tdap as part of the catch-up series require 1 additional dose at 11-12 for 7th grade entry.

Polio: A minimum of 4 weeks between doses required with 6 months between last two doses. *OPV: Only trivalent OPV counts as valid. Monovalent or bivalent OPV are not valid. All doses given after 4/1/16 are assumed to be mono or bivalent.

MMR: Required 2nd dose should be given on/after 4th birthday. However, dose 2 may be given earlier with at least 4 weeks between dose 1 and 2.

Hib: If series started <12 months, 3 doses req'd with at least 1 dose on/after 1st Bday. Two doses req'd if dose 1 was given at 12-14 months. One dose of Hib given between 16-59 months is sufficient. Not recommended ≥60 months.

Hep B: Dose 2 a minimum of 4 weeks after dose 1; dose 3 at least 16 weeks after dose 1 and at least 8 weeks after dose 2. Infants currently receiving primary series, final dose should be administered no earlier than age 24 weeks.

PCV: Administer a series of PCV13 vaccine at ages 2, 4, 6 months with a booster at age 12-15 months. Catch-up: Administer one dose of PCV13 to all healthy children 12-59 months who are not completely vaccinated for their age; children >60 months, no doses required.

Varicella: For children aged 12 months-12 years, the minimum interval between the two doses is 3 months. However, if dose 2 was administered ≥28 days after dose 1, dose 2 is considered valid and need not be repeated. For children ≥13 years, the recommended minimum interval is 4 wks. **Required for proof of varicella immunity:**

- For K - 7th graders: Receipt of vaccine; titer or laboratory confirmed diagnosis is required as proof of prior disease.
- For 8th - 12th graders: Receipt of vaccine, written proof of immunity by a physician/health care provider or laboratory titer is required.
- For all newly diagnosed varicella cases: Laboratory confirmation of disease is required.

Hep A: One dose required by 15 months; 2 doses required at 48 mos. with at least 6 months between doses.

Recommended # doses for adult students 19+ in secondary school

Vaccine	doses
Tetanus/Diphtheria/Pertussis (Tdap)	1
Measles/Mumps/Rubella (MMR)	2
Varicella (VAR)	2

Resources

CDC Immunization Schedule has detailed footnotes and catch-up schedule <https://www.cdc.gov/vaccines/schedules/hcp/imz/c-hild-adolescent.html>

NM Immunization Protocol <https://nmhealth.org/publication/view/regulation/531/>

NMSIIS <https://nmsiis.health.state.nm.us>

NM School Health Manual <http://www.nmschoolhealthmanual.org>

*Minimum age 6 wks. Age-appropriate flu & COVID vaccines are recommended for all. Changes from last year's requirements are highlighted for ease of use.

Note: Any vaccine administered ≤4 days prior to minimum interval or age is valid. All students enrolled in designated grades are expected to meet requirements.